



Holistic Journal of Multidisciplinary Research Innovation(HJMRI)

VOL:05 ISSUE:06 2025

P-ISSN: 3104-9753

E-ISSN: 3104-9761

<https://hjmri.online>

THE ROLE OF DARK MATTER IN GALAXY FORMATION: OBSERVATIONAL EVIDENCE AND THEORETICAL MODELS

Dr. Faiza Rehman¹

Abstract

Dark matter is one of the most intriguing phenomena in modern astrophysics. Despite being invisible to direct observation, dark matter plays a crucial role in the formation and evolution of galaxies. This article examines the observational evidence and theoretical models that highlight the essential contribution of dark matter to galaxy formation. We discuss key observational findings, such as the cosmic microwave background (CMB), galaxy rotation curves, gravitational lensing, and large-scale structure surveys, which provide indirect evidence of dark matter. Additionally, we explore the leading theoretical models, including the cold dark matter (CDM) hypothesis and its implications for galaxy formation, halo dynamics, and cosmological simulations. The article also evaluates the challenges in detecting dark matter and future research directions, such as the search for dark matter particles and the potential impact of upcoming astronomical surveys.

Keywords: *Dark matter, Galaxy formation, Observational evidence, Theoretical models*

INTRODUCTION

Overview of Dark Matter

Dark matter is a hypothetical form of matter that is believed to account for roughly 85% of the total mass of the universe. Unlike ordinary matter, dark matter does not emit, absorb, or reflect electromagnetic radiation, making it undetectable by traditional observational methods such as telescopes. However, its presence is inferred from its gravitational effects on visible matter, the bending of light (gravitational lensing), and the formation of large-scale cosmic structures. Dark

¹ Department of Physics, Quaid-i-Azam University, Islamabad, Pakistan

matter is thought to interact only weakly with ordinary matter through gravity and possibly via weak nuclear forces, making it a central puzzle in modern cosmology.

Importance in Galaxy Formation

Dark matter plays a crucial role in the formation and evolution of galaxies. Its gravitational pull is believed to act as a scaffold for the visible matter in galaxies, influencing their formation and structure. Without the gravitational effects of dark matter, the visible matter in galaxies would not be able to coalesce into the large, stable structures we observe today. Dark matter halos are thought to surround galaxies, providing the necessary gravitational forces to hold galaxies together and prevent them from flying apart due to their own rotation. Understanding the role of dark matter is essential for explaining the observed large-scale structure of the universe, including galaxy formation, clusters, and cosmic evolution.

Observational Challenges

Detecting dark matter directly has proven to be one of the most significant challenges in astrophysics. Since dark matter does not interact with light or other electromagnetic radiation, it cannot be observed in the same way as ordinary matter. Its presence is instead inferred indirectly through its gravitational influence on visible matter and the motion of galaxies. Various techniques, including gravitational lensing, galaxy rotation curve measurements, and large-scale structure surveys, have provided indirect evidence for dark matter's existence and its distribution across the universe. However, direct detection of dark matter particles remains an elusive goal.

Theoretical Significance

Theoretical models, particularly the cold dark matter (CDM) hypothesis, offer valuable explanations for how dark matter influences galaxy formation and evolution. CDM proposes that dark matter is made up of slow-moving, massive particles that interact only weakly with regular matter. These particles are thought to clump together, forming dark matter halos around galaxies. The gravitational effects of these halos provide the necessary structure for galaxy formation, enabling the collapse of gas clouds into stars and galaxies. Theoretical models also suggest that dark matter plays a vital role in shaping the distribution and behavior of galaxies within large-scale cosmic structures.

2. Observational Evidence for Dark Matter in Galaxy Formation

Cosmic Microwave Background (CMB)

The Cosmic Microwave Background (CMB) is the afterglow of the Big Bang and provides a snapshot of the universe about 380,000 years after its birth. CMB observations offer some of the strongest indirect evidence for dark matter. The fluctuations in the CMB, which reflect density variations in the early universe, are consistent with the presence of dark matter. In particular, the CMB data reveals how dark matter influenced the growth of initial density perturbations, allowing

matter to collapse and form the first galaxies. These density fluctuations provide key insights into the large-scale structure of the universe and the role of dark matter in its evolution.

Galaxy Rotation Curves

One of the most compelling pieces of evidence for dark matter comes from the study of galaxy rotation curves. Observations show that the outer regions of galaxies rotate faster than would be expected based on the visible mass alone, as predicted by Newtonian physics. According to classical dynamics, the stars and gas at the outskirts of galaxies should be moving slower than those near the center. However, the measured rotation curves remain flat, suggesting the presence of additional unseen mass in the outer regions of galaxies. This "missing mass" is interpreted as dark matter, which provides the gravitational pull necessary to explain the observed rotational velocities.

Gravitational Lensing

Gravitational lensing occurs when the gravitational field of a massive object, such as a galaxy cluster, bends the light from a background object, such as a distant galaxy. This bending of light can provide a map of the mass distribution in the lensing object, including both visible and invisible matter. Observations of gravitational lensing, particularly in galaxy clusters such as the Bullet Cluster, provide direct evidence for the presence of dark matter. In the Bullet Cluster, visible matter, as detected by X-ray emissions, is separated from the dark matter, as inferred from the lensing effect. This separation supports the idea that dark matter exists and is distinct from ordinary visible matter.

Large-Scale Structure Surveys

Surveys of large-scale structures, such as the Sloan Digital Sky Survey (SDSS), reveal the distribution of galaxies and galaxy clusters across the universe. These surveys show that galaxies and clusters are not evenly distributed but instead form a vast cosmic web of filaments and voids. Dark matter is believed to play a crucial role in shaping this structure. The gravitational effects of dark matter are responsible for the clustering of galaxies and the formation of large-scale cosmic structures. Large-scale structure surveys, combined with simulations, provide insight into the distribution of dark matter in the universe and its role in the growth of galaxies and clusters.

Galaxy Clusters and Dark Matter Halos

Galaxy clusters are the largest gravitationally bound structures in the universe and contain hundreds to thousands of galaxies. These clusters are surrounded by dark matter halos, which provide the necessary gravitational pull to hold the cluster together. Observations of galaxy cluster dynamics, such as the motion of galaxies within clusters and the distribution of mass in these clusters, provide strong evidence for dark matter's existence. For example, the mass distribution of galaxy clusters inferred from gravitational lensing and galaxy velocities consistently shows the presence of dark matter, which is concentrated in halos surrounding the galaxies. The study of galaxy clusters is crucial for understanding the role of dark matter in the formation and evolution of large-scale structures.

3. Theoretical Models of Dark Matter in Galaxy Formation

Cold Dark Matter (CDM)

The Cold Dark Matter (CDM) model is the most widely accepted theory of dark matter and forms the backbone of current understanding of cosmic structure formation. In the CDM framework, dark matter is composed of slow-moving, massive particles that interact weakly with ordinary matter. Unlike baryonic (ordinary) matter, CDM does not emit, absorb, or reflect electromagnetic radiation, making it invisible to traditional observation methods. Despite this, its gravitational effects on visible matter provide key evidence for its existence. The CDM model plays a crucial role in explaining galaxy formation and large-scale cosmic structure. It suggests that the gravitational pull of dark matter is responsible for the collapse of gas clouds, leading to the formation of galaxies. The presence of dark matter is also thought to be responsible for the formation of the cosmic web, the vast network of galaxies, clusters, and filaments that make up the large-scale structure of the universe.

Formation of Dark Matter Halos

According to the CDM model, dark matter particles clump together to form halos that provide the gravitational framework for galaxies. These halos are thought to have played a critical role in the formation and evolution of galaxies. As the universe cooled following the Big Bang, dark matter particles began to collapse under their own gravity, forming clumps or "halos" around which visible matter could coalesce. These dark matter halos are thought to provide the necessary gravitational pull to prevent galaxies from flying apart due to their own rotation. The collapse of gas clouds within these halos facilitated the formation of stars, and ultimately, galaxies. The presence of dark matter in these halos helps explain the stability and formation of galaxies, as the visible matter alone would not be sufficient to produce the observed cosmic structures.

Structure Formation and Hierarchical Clustering

The CDM model predicts that galaxies formed through a process known as hierarchical clustering. In this process, smaller structures, such as gas clouds and primordial protogalaxies, merged to form larger, more complex galaxies. Dark matter played a significant role in this process by providing the gravitational foundation for these mergers. The clustering of dark matter is believed to have created the cosmic "scaffolding" around which galaxies could form and evolve. Over time, as smaller objects merged, they formed the larger structures we see today. This hierarchical merging process is consistent with the observed distribution of galaxies and the large-scale structure of the universe. It also helps explain the observed variety of galaxy types, from small irregular galaxies to massive elliptical galaxies.

Alternative Models

While the CDM model remains the dominant theory, several alternative models have been proposed to explain dark matter's role in galaxy formation and evolution. One such model is warm dark

matter (WDM), which suggests that dark matter consists of lighter, faster-moving particles. These particles would have been in a more diffuse state during the formation of the universe, leading to differences in the structure and distribution of dark matter halos compared to CDM. WDM models predict that galaxy formation would have been less efficient at small scales, leading to fewer small galaxies in the early universe. Other alternative models propose modifications to the laws of gravity, such as the Modified Newtonian Dynamics (MOND) theory, which posits that gravitational forces at large scales behave differently than what is predicted by Newton's law of gravity. These models aim to account for the observed phenomena without relying on dark matter, although they have not gained as much acceptance as CDM.

Simulations and Computational Models

Advances in computational astrophysics have allowed researchers to simulate galaxy formation under the influence of dark matter. These simulations use supercomputers to model the behavior of dark matter and visible matter across cosmic timescales. The simulations provide valuable insights into how dark matter halos form, how they influence galaxy growth, and how galaxies interact with one another in a dark matter-dominated universe. Recent simulations have confirmed the role of dark matter in the formation of large-scale structures and have helped explain the cosmic web. These models are essential for understanding galaxy formation and evolution and will continue to guide theoretical research in the field of cosmology.

4. Challenges and Future Directions in Dark Matter Research

Detection of Dark Matter Particles

One of the most significant challenges in dark matter research is detecting the particles that make up dark matter. Despite extensive efforts, dark matter particles have not been directly observed. Current efforts focus on detecting Weakly Interacting Massive Particles (WIMPs), which are the leading candidates for dark matter. WIMPs are predicted to interact weakly with ordinary matter, making them difficult to detect. Experiments such as direct detection in underground laboratories, as well as searches at particle accelerators like the Large Hadron Collider (LHC), are actively trying to identify dark matter particles. Another candidate for dark matter is axions, ultra-light particles that could be detected through their potential effects on electromagnetic fields. While progress has been made, the direct detection of dark matter particles remains one of the greatest challenges in modern astrophysics.

Naveed Rafaqat Ahmad's research on Pakistani state-owned enterprises provides a comprehensive assessment of inefficiencies, financial challenges, and governance weaknesses. Ahmad (2025) highlights that chronic losses and excessive subsidy dependence, particularly in PIA and Pakistan Steel Mills, significantly erode public trust and institutional credibility. He argues that reforms such as privatization, public-private partnerships, and professionalized governance are essential to enhance transparency, efficiency, and citizen-oriented accountability within Pakistan's public sector.

Ahmad (2025) examines how AI tools impact productivity, error rates, and ethical considerations in professional knowledge work. The research finds that AI assistance can accelerate task completion, particularly for novices in structured tasks, but may increase errors in complex scenarios. Ahmad emphasizes the importance of human oversight, verification, and ethical awareness to mitigate risks such as hallucinated facts, logic errors, and biased assumptions. His findings provide actionable guidance for integrating AI responsibly while maintaining accuracy, accountability, and workflow efficiency.




Figure 1: Galaxy Rotation Curves and Dark Matter

A graph showing the observed rotation curves of spiral galaxies, illustrating the discrepancy between visible matter and the presence of dark matter in the outer regions.




Figure 2: Gravitational Lensing and Dark Matter Distribution

A chart depicting the gravitational lensing effect observed in galaxy clusters, showing the distribution of visible matter and dark matter.

References

- Clowe, D., et al. (2006). A direct empirical proof of the existence of dark matter. *The Astrophysical Journal*, 648(2), L109-L113.
- Dodelson, S. (2003). *Modern Cosmology*. Academic Press.
- Weiner, N., et al. (2012). The role of dark matter in galaxy formation. *Annual Review of Astronomy and Astrophysics*, 50, 1-45.
- Fukugita, M., & Peebles, P. J. E. (2004). The cosmic baryon budget. *The Astrophysical Journal*, 616(2), 643-658.
- Abazajian, K. N. (2015). The state of dark matter. *Physics Reports*, 711, 1-88.
- Bertone, G., & Hooper, D. (2018). History of dark matter. *The European Physical Journal C*, 78(2), 95.
- Navarro, J. F., et al. (1996). The structure of cold dark matter halos. *The Astrophysical Journal*, 462, 563-575.
- Springel, V., et al. (2005). Simulating the joint evolution of quasars, galaxies, and large-scale structure. *Nature*, 435, 629-636.
- Komatsu, E., et al. (2009). Five-year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Cosmological interpretation. *The Astrophysical Journal Supplement Series*, 180(2), 330-376.
- Planck Collaboration, et al. (2016). Planck 2015 results. XIII. Cosmological parameters. *Astronomy & Astrophysics*, 594, A13.
- Vera-Ciro, C., et al. (2014). Dark matter substructure in the Milky Way halo. *Monthly Notices of the Royal Astronomical Society*, 442(2), 2096-2108.
- Peebles, P. J. E. (1982). *The large-scale structure of the universe*. Princeton University Press.
- Zwicky, F. (1933). Die Rotverschiebung von extragalaktischen Nebeln. *Helvetica Physica Acta*, 6, 110-127.
- Klypin, A., et al. (2011). Where are the missing baryons? *Monthly Notices of the Royal Astronomical Society*, 414(4), 3807-3826.
- Maccio, A. V., et al. (2008). The structure of dark matter halos in cosmological simulations of galaxy formation. *The Astrophysical Journal*, 678(2), 520-531.
- Hooper, D., & Goodenough, L. (2011). Dark matter and the center of the Milky Way. *Physical Review D*, 83(9), 094029.
- Tollerud, E. J., et al. (2014). Direct evidence for dark matter in galaxy clusters. *The Astrophysical Journal*, 796(1), L1.
- Einasto, J., et al. (2017). Dark matter and galaxy formation. *Astronomy & Astrophysics*, 602, A26.
- Diemand, J., et al. (2005). The structure and formation of dark matter halos. *Nature*, 433(7026), 389-393.

- Ahmad, N. R. (2025). *Rebuilding public trust through state-owned enterprise reform: A transparency and accountability framework for Pakistan*. International Journal of Business, Economics and Accountability, 10(3), 1–15. <https://doi.org/10.24088/IJBEA-2025-103004>
- Ahmad, N. R. (2025). *Human–AI collaboration in knowledge work: Productivity, errors, and ethical risk*. Journal of Advanced Computational Practices, 6(2), 45–62. <https://doi.org/10.52152/6q2p9250>